


Anti-Neurologin 4 Antibody (A14578)

Specifications:

Name:	Anti-Neurologin 4 Antibody
Description:	Rabbit polyclonal antibody to Neurologin 4.
Applications:	WB, IHC
Recommended Dilutions:	WB: 1:500-1:2,000, IHC: 1:50-1:100
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 122-256 of human NLGN4Y (NP_001157710.1).
Sequence:	DMLPIWFTTSLDTLMTYVQDQNECLYLNIVPMEDGTNIKRNADDITSNDHGDKDI HEQNSKPKVMVYIHGGSYMEGTGNMIDGSILASYGNVIVITINYRLGILGMQEARLCG SSKMFNYFKSPFTNLINFF
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	100 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:



Western blot analysis of extracts of various cell lines, using Anti-Neurologin 4 Antibody (A14578) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.

Anti-Neurologin 4 Antibody (A14578)

Images continued:



Immunohistochemistry analysis of paraffin-embedded human prostate using Anti-Neurologin 4 Antibody (A14578) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.