


Anti-HDAC8 Antibody (A15002)

Specifications:


Name:	Anti-HDAC8 Antibody
Description:	Rabbit polyclonal antibody to HDAC8.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:500-1:1,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-377 of human HDAC8 (NP_060956.1).
Sequence:	MEEPEEPADSGQSLVPVYIYSPEYVSMCDSLAKIPKRASMVHSLIEAYALHKQMRIVK PKVASMEEMATFHTDAYLQHLQKVSQEGDDDHDPDSIEYGLGYDCPATEGIFDYAAAIG GATITAAQCLIDGMCKVAINWSGGWHHAKKDEASGFCYLNDVAVLGILRLRRKFERILY VDLDLHHGDGVEDAFSFTSKVMTVSLHKFSPGFFPGTGDVSDVGLGKGRYYSVNVPIQ DGIQDEKYYQICESVLKEVYQAFNPKAVVLQLGADTIAGDPMCSFNMTVPVIGIGKCLKY ILQWQLATLILGGGGYNLANTARCWTYLTGVILGKTLSSSEIPDHEFFTA YGPDYVLEI TPSCRPRNEPHRIQQILNYIKGNLKHVV
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	45 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-HDAC8 Antibody (A15002)


Images:




Western blot analysis of extracts from normal (control) and HDAC8 knockout (KO) HeLa cells, using Anti-HDAC8 Antibody (A15002) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.



Western blot analysis of extracts of C6 cells, using Anti-HDAC8 Antibody (A15002) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded rat testis using Anti-HDAC8 Antibody (A15002) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat spinal cord using Anti-HDAC8 Antibody (A15002) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.