

Anti-SSRP1 Antibody (A15218)

Specifications:

Name: Anti-SSRP1 Antibody

Description: Rabbit polyclonal antibody to SSRP1.

Applications: WB, IHC, ICC/IF

Recommended Dilutions: WB: 1:100-1:500, IHC: 1:50-1:200, ICC/IF: 1:10-1:100

Reactivity: Human, Mouse

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of

human SSRP1 (NP_003137.1).

Sequence: MAETLEFNDVYQEVKGSMNDGRLRLSRQGIIFKNSKTGKVDNIQAGELTEGIWRRVAL

GHGLKLLTKNGHVYKYDGFRESEFEKLSDFFKTHYRLELMEKDLCVKGWNWGTVKFGG QLLSFDIGDQPVFEIPLSNVSQCTTGKNEVTLEFHQNDDAEVSLMEVRFYVPPTQEDG VDPVEAFAQNVLSKADVIQATGDAICIFRELQCLTPRGRYDIRIYPTFLHLHGKTFDY KIPYTTVLRLFLLPHKDQRQMFFVISLDPPIKQGQTRYHFLILLFSKDEDISLTLNMN

EEEVEKRFEG

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 90 kDa

Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300.


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




Anti-SSRP1 Antibody (A15218)


Images:



Western blot analysis of extracts of Mouse brain, using Anti-SSRP1 Antibody (A15218) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 5s.



Immunohistochemistry analysis of paraffin-embedded mouse liver using Anti-SSRP1 Antibody (A15218) at a dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of U2OS cells using Anti-SSRP1 Antibody (A15218).