


## Anti-LOX Antibody (A15846)

### Specifications:


Name:	Anti-LOX Antibody
Description:	Rabbit polyclonal antibody to LOX.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:100-1:500, IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-168 of human LOX (NP_002308.2).
Sequence:	MRFAWTVLLLGLPLQLCALVHCAPPAAGQQQPPREPPAAPGAWRQQIQWENNGQVFSLL SLGSQYQPQRRRDPGAAVPGAANASAAQQRTPILLIRDNRATAARTRTAGSSGVTAGR PRPTARHWFQAGYSTSRAREAGASRAENQTAPGEVPALSNLRPPSRVDGMVG
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	47 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-LOX Antibody (A15846)


### Images:




Western blot analysis of extracts of various cell lines, using Anti-LOX Antibody (A15846) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 90s.



Immunofluorescence analysis of BALB-3T3 cells using Anti-LOX Antibody (A15846) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of C6 cells using Anti-LOX Antibody (A15846) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of U-2 OS cells using Anti-LOX Antibody (A15846) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).