


Anti-Serine / threonine-protein kinase 4 / MST-1 Antibody (A16052)

Specifications:


Name:	Anti-Serine / threonine-protein kinase 4 / MST-1 Antibody
Description:	Rabbit polyclonal antibody to Serine / threonine-protein kinase 4 / MST-1.
Applications:	WB, IHC, ICC/IF, IP
Recommended Dilutions:	WB: 1:500-1:1,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200, IP: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 303-487 of human MST1/STK4 (NP_006273.1).
Sequence:	KRQESQQREVDQDDEENSEEDEMDSGTMVRAVGDEMGTVRVASTMTDGANTMIEHDDT LPSQLGTMVINAEEEEEGTMKRRDETMQPAKPSFLEYFEQKEKENQINSFGKSVPGP LKNSSDWKIPQDGDYEFKSWTVEDLQKRLLALDPMMEQEIEEIRQKYQSKRQPILDA IEAKKRRQQNF
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	56 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-Serine / threonine-protein kinase 4 / MST-1 Antibody (A16052)


Images:



Western blot analysis of extracts of various cells, using Anti-Serine / threonine-protein kinase 4 / MST-1 Antibody (A16052) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 15s.



Western blot analysis of extracts from normal (control) and MST1/STK4 knockout (KO) 293T cells, using Anti-Serine / threonine-protein kinase 4 / MST-1 Antibody (A16052) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.



Immunoprecipitation analysis of 100µg extracts of HepG2 cells using 3µg of Anti-Serine / threonine-protein kinase 4 / MST-1 Antibody (A16052). This Western blot was performed on the immunoprecipitate using Anti-Serine / threonine-protein kinase 4 / MST-1 Antibody (A16052) at a dilution of 1:1000.