


Anti-RIP3 Antibody (A17064)

Specifications:

Name:	Anti-RIP3 Antibody
Description:	Rabbit polyclonal antibody to RIP3.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:500-1:2,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human RIP3 (NP_006862.2).
Sequence:	GSLSGLLQSQCPRPWPLLCRLLKEVVLGMFYLDQNPVLLHRDLKPSNVLLDPELHVK LADFGLSTFQGGSQSGTGSGEPGGTLGYLAPELNVNVRKAST
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	46 - 62 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:




Western blot analysis of various lysates, using Anti-RIP3 Antibody (A17064) at 1:3,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.

Anti-RIP3 Antibody (A17064)


Images continued:



Immunohistochemistry analysis of paraffin-embedded human liver cancer using Anti-RIP3 Antibody (A17064) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse pancreas using Anti-RIP3 Antibody (A17064) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of C6 cells using Anti-RIP3 Antibody (A17064) at a dilution of 1:50 (40x lens). DAPI was used to stain the cell nuclei (blue).