


Anti-PRDM2 Antibody (A17292)

Specifications:

Name:	Anti-PRDM2 Antibody
Description:	Rabbit polyclonal antibody to PRDM2.
Applications:	WB, IHC
Recommended Dilutions:	WB: 1:500-1:2000, IHC: 1:50-1:200
Reactivity:	Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-240 of human PRDM2 (NP_036363.2).
Sequence:	MNQNTTEPVAATETLAEVPEHVLRLGLPEEVRLFPSAVDKTRIGVWATKPIKGGKFGP FVGDKKKRSQVKNNVYMWEVYYPNLGWMCIDATDPEKGNWLRVYNWACSGEEQNLFP EINRAIYYKTLKPIAPGEELLVWYNGEDNPEIAAAIEEERASARSKRSSPKSRKGKKK SQENKNKGNKIQDIQLKTSEPDFTSANMRDSAEGPKEDEEKPSASALEQPATLQEVAS QEVPPELA
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	250kDa
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.30, with 0.02% Sodium Azide and 50% Glycerol.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-PRDM2 Antibody (A17292)

Images:



Western blot analysis of extracts of rat heart, using Anti-PRDM2 Antibody (A8565) at 1:1,000 dilution.


Secondary antibody: Goat Anti-Rabbit IgG (H+L) (HRP) (AS014) at 1:10,000 dilution.

Lysates / proteins: 25µg per lane.


Blocking buffer: 3% non-fat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 90s.



Immunohistochemistry of paraffin-embedded rat lung using Anti-PRDM2 Antibody (A8565) at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded rat liver using Anti-PRDM2 Antibody (A8565) at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded rat testis using Anti-PRDM2 Antibody (A8565) at dilution of 1:100 (40x lens).