


## Anti-TIF1 alpha Antibody [PCRP-TRIM24-1B12] (A250442)

### Specifications:


Name:	Anti-TIF1 alpha Antibody [PCRP-TRIM24-1B12]
Description:	Mouse monoclonal [PCRP-TRIM24-1B12] antibody to TIF1 alpha.
Specificity:	TIF1 homolog (designated bonus) has been identified in Drosophila and is associated with several genes that are implicated in the ecdysone pathway, a nuclear hormone receptor pathway required throughout Drosophila development, suggesting a conserved functional role for the protein throughout the course of evolution.
Applications:	Flow Cytometry, IF
Recommended Dilutions:	Flow Cytometry: 1-2 µg/million cells, IF: 1-2 µg/ml
Reactivity:	Human
Immunogen:	Recombinant full-length human TRIM24 protein.
Host:	Mouse
Clonality:	Monoclonal
Clone ID:	PCRP-TRIM24-1B12
Isotype:	IgG1
Conjugate:	Unconjugated
Purification:	Protein A/G chromatography.
Concentration:	200 µg/ml
Product Form:	Liquid
Formulation:	Supplied in 10mM Phosphate Buffered Saline with 0.05% BSA and 0.05% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
General Notes:	This monoclonal antibody is also available in a different formulation without BSA and Sodium Azide - Anti-TIF1 alpha Antibody [PCRP-TRIM24-1B12] - BSA and Azide free (A253622).
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

# Anti-TIF1 alpha Antibody [PCRP-TRIM24-1B12] (A250442)


## Images:



Flow cytometric analysis of PFA fixed HeLa cells using Anti-TIF1 alpha Antibody [PCRP-TRIM24-1B12] followed by Goat Anti-Mouse IgG (CF® 488) (Blue). Isotype Control (Red).



Immunofluorescent analysis of PFA fixed MCF-7 cells stained with Anti-TIF1 alpha Antibody [PCRP-TRIM24-1B12] followed by Goat Anti-Mouse IgG (CF® 488) (Green). Counterstain is Phalloidin.



Analysis of protein array containing more than 19,000 full-length human proteins using Anti-TIF1 alpha Antibody [PCRP-TRIM24-1B12]. Z-Score and S-Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target; a MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.