


## Anti-HIC2 Antibody [PCRP-HIC2-1B1] (A277608)

### Specifications:


Name:	Anti-HIC2 Antibody [PCRP-HIC2-1B1]
Description:	Mouse monoclonal [PCRP-HIC2-1B1] antibody to HIC2.
Specificity:	HIC2 (HIC ZBTB Transcriptional Repressor 2) is a Protein Coding gene. Diseases associated with HIC2 include Orofaciodigital Syndrome X and Simpson-Golabi-Behmel Syndrome, Type 1. HIC2 contains 5 C2H2-type zinc fingers and 1 BTB (POZ) domain. It belongs to the krueppel C2H2-type zinc-finger protein family, Hic subfamily and is a transcriptional repressor. It is a transcription activator of SIRT1.
Applications:	ELISA, IP, Flow Cytometry, IF
Recommended Dilutions:	IP: 1-2µg / 100-500µg proteins, Flow Cytometry: 1-2 µg/million cells, IF: 1-2 µg/ml
Reactivity:	Human
Immunogen:	Recombinant full-length human HIC2 protein.
Host:	Mouse
Clonality:	Monoclonal
Clone ID:	PCRP-HIC2-1B1
Isotype:	IgG2a
Conjugate:	Unconjugated
Purification:	Protein A/G chromatography.
Concentration:	200 µg/ml
Product Form:	Liquid
Formulation:	Supplied in 10mM Phosphate Buffered Saline with 0.05% BSA and 0.05% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
General Notes:	This monoclonal antibody is also available in a different formulation without BSA and Sodium Azide - Anti-HIC2 Antibody [PCRP-HIC2-1B1] - BSA and Azide free (A278196).
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-HIC2 Antibody [PCRP-HIC2-1B1] (A277608)


### Images:



Flow cytometric analysis of PFA-fixed HeLa cells using Anti-HIC2 Antibody [PCRP-HIC2-1B1] followed by Goat Anti-Mouse IgG (CF® 488) (Blue). Isotype Control (Red).




SDS-PAGE analysis of Anti-HIC2 Antibody [PCRP-HIC2-1B1] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.



Immunofluorescent analysis of PFA-fixed HeLa cells stained with Anti-HIC2 Antibody [PCRP-HIC2-1B1] followed by Goat Anti-Mouse IgG (CF® 488) (Green). CF® 640A Phalloidin (Red).

## Anti-HIC2 Antibody [PCRP-HIC2-1B1] (A277608)

Images continued:



Analysis of protein array containing more than 19,000 full-length human proteins using Anti-HIC2 Antibody [PCRP-HIC2-1B1]. Z-Score and S-Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target; a MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.