


Anti-Galectin 3 Antibody [LGALS3/4792] - BSA and Azide free (A278279)

Specifications:


Name:	Anti-Galectin 3 Antibody [LGALS3/4792] - BSA and Azide free
Description:	Mouse monoclonal [LGALS3/4792] antibody to Galectin 3.
Specificity:	Galectin-3 is a member of the beta-galactosidase-binding lectin family. It is associated with cell growth, adhesion, inflammation, mRNA processing, and apoptosis. Aberrant expression of Galectin-3 is related to malignant transformation and metastasis in carcinomas of the breast, colon and thyroid. Galectin-3 reactivity can be seen in the nucleus of neutrophils, vascular endothelium, carcinomas of the colon, breast, and thyroid. Galectin-3 may be useful in the differentiation of benign and malignant thyroid neoplasms. Galectin-3 may also be useful in the identification of certain liver disorders.
Applications:	IHC-P
Recommended Dilutions:	IHC-P: 1-2 µg/ml
Reactivity:	Human
Immunogen:	Synthetic peptide corresponding to the C terminus, around amino acids 150-200, of human Galectin 3 protein. The exact sequence is proprietary.
Host:	Mouse
Clonality:	Monoclonal
Clone ID:	LGALS3/4792
Isotype:	IgG1
Light Chains:	kappa
Conjugate:	Unconjugated
Purification:	Protein A/G chromatography.
Concentration:	1 mg/ml
Product Form:	Liquid
Formulation:	Supplied in 10mM Phosphate Buffered Saline; without Sodium Azide and carrier free.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
General Notes:	This monoclonal antibody is also available in a different formulation with BSA and Sodium Azide - Anti-Galectin 3 Antibody [LGALS3/4792] (A277691).
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-Galectin 3 Antibody [LGALS3/4792] - BSA and Azide free (A278279)


Images:



Immunohistochemical analysis of formalin-fixed, paraffin-embedded human thyroid carcinoma tissue using Anti-Galectin 3 Antibody [LGALS3/4792].




Immunohistochemical analysis of formalin-fixed, paraffin-embedded human thyroid carcinoma tissue using Anti-Galectin 3 Antibody [LGALS3/4792].



SDS-PAGE analysis of Anti-Galectin 3 Antibody [LGALS3/4792] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.

Anti-Galectin 3 Antibody [LGALS3/4792] - BSA and Azide free (A278279)

Images continued:



Analysis of protein array containing more than 19,000 full-length human proteins using Anti-Galectin 3 Antibody [LGALS3/4792]. Z-Score and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target; a MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.