


Anti-NMDAR1 Antibody [S308-48] (A304953)

Specifications:

Name:	Anti-NMDAR1 Antibody [S308-48]
Description:	Mouse monoclonal [S308-48] antibody to NMDAR1.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:1,000, IHC: 1:1,000, ICC/IF: 1:100
Reactivity:	Human, Mouse, Rat
Immunogen:	Fusion protein corresponding to the extracellular N-terminus, amino acids 42-361, of rat NR1 protein.
Host:	Mouse
Clonality:	Monoclonal
Clone ID:	S308-48
Isotype:	IgG1
Conjugate:	Unconjugated
Purification:	Protein G purification.
Concentration:	1 mg/ml
Molecular Weight:	~105 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.4, with 50% Glycerol and 0.09% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:




Western blot analysis of rat brain membrane lysate showing detection of NMDAR1 NMDA receptor protein using Anti-NMDAR1 Antibody [S308-48] (A304953) at 1:1,000.

Anti-NMDAR1 Antibody [S308-48] (A304953)

Images continued:



Immunocytochemistry/Immunofluorescence analysis of human neuroblastoma cell line (SK-N-BE, fixed in 4% formaldehyde for 15 min at room temperature, using Anti-NMDAR1 Antibody [S308-48] (A304953), at 1:100 for 60 minutes at room temperature. The secondary antibody used was Goat Anti-Mouse ATTO 488 at 1:100 for 60 minutes at room temperature. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5,000 for 60min room temperature, 5min room temperature. Localization: Cell Membrane. Magnification: 60X. (A) DAPI (blue) nuclear stain. (B) Phalloidin Texas Red F-Actin stain. (C) GluN1/NR1 Antibody. (D) Composite.



Immunocytochemistry/Immunofluorescence analysis of human neuroblastoma cells (SH-SY5Y), fixed in 4% PFA for 15 min, using Anti-NMDAR1 Antibody [S308-48] (A304953), at 1:50 for overnight at 4°C with slow rocking. The secondary antibody used was AlexaFluor 488 at 1:1,000 for 1 hour at room temperature. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain; Hoechst (blue) nuclear stain at 1:800, 1.6mM for 20 minutes at room temperature. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) GluN1/NR1 Antibody (D) Composite.