


Anti-iNOS Antibody [ARC0832] (A305283)

Specifications:


Name:	Anti-iNOS Antibody [ARC0832]
Description:	Rabbit monoclonal [ARC0832] antibody to iNOS.
Applications:	WB, ICC/IF
Recommended Dilutions:	WB: 1:500-1:1,000, ICC/IF: 1:50-1:200
Reactivity:	Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 2-200 of human iNOS (P35228).
Sequence:	ACPWKFLFKTKFHQYAMNGEKDINNNVEKAPCATSSPVTQDDLQYHNLSKQQNESPQP LVETGKKSPESLVKLDATPLSSPRHVRIKNWGSMTFQDTLHHKAKGILTCRSKSLG SIMTPKSLTRGPRDKPTPPDELLPQAIEFVNQYYGSFKEAKIEEHLARVEAVTKEIET TGTYQLTGDELIFATKQAWRNAPRC
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC0832
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	130 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-iNOS Antibody [ARC0832] (A305283)

Images:



Western blot analysis of extracts of RAW264.7 cells, using Anti-iNOS Antibody [ARC0832] (A305283) at 1:1,000 dilution. RAW264.7 cells were treated by LPS (1 $\mu\text{g}/\text{ml}$) at 37°C for 8 hours. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 180s.



Immunofluorescence analysis of Raw264.7 and Raw264.7 treated with LPS cells using Anti-iNOS Antibody [ARC0832] (A305283) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).