


Anti-Cytokeratin 18 Antibody [ARC2303] (A305288)

Specifications:


Name:	Anti-Cytokeratin 18 Antibody [ARC2303]
Description:	Rabbit monoclonal [ARC2303] antibody to Cytokeratin 18.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:1,000-1:3,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 250-430 of human Cytokeratin 18 (KRT18) (P05783).
Sequence:	ADIRAQYDELARKNREELDKYWSQQIEESTTVVTTQSAEVGAAETTLTELRRTVQSLE IDLDSMRNLKASLENSLREVEARYALQMEQLNGILLHLESELAQTRAEGQRQAQEYEA LLNIKVKLEAEIATYRRILLEDGEDFNLGDALDSSNSMQTIQKTTTRRIVDGKVVSETN DTKVLRH
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC2303
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	48 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-Cytokeratin 18 Antibody [ARC2303] (A305288)


Images:




Western blot analysis of extracts of various cell lines, using Anti-Cytokeratin 18 Antibody [ARC2303] (A305288) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.



Immunohistochemistry analysis of paraffin-embedded rat liver using Anti-Cytokeratin 18 Antibody [ARC2303] (A305288) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human liver cancer using Anti-Cytokeratin 18 Antibody [ARC2303] (A305288) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse stomach using Anti-Cytokeratin 18 Antibody [ARC2303] (A305288) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.