

## Anti-PIAS1 Antibody [ARC1116] (A305639)

## Specifications:

Name: Anti-PIAS1 Antibody [ARC1116]

Description: Rabbit monoclonal [ARC1116] antibody to PIAS1.

Applications: WB

Recommended Dilutions: WB: 1:500-1:2,000

Reactivity: Human, Mouse

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 3-101 of

human PIAS1 (075925).

Sequence: DSAELKQMVMSLRVSELQVLLGYAGRNKHGRKHELLTKALHLLKAGCSPAVQMKIKEL

YRRRFPQKIMTPADLSIPNVHSSPMPATLSPSTIPQLTYDG

Host: Rabbit

Clonality: Monoclonal

Clone ID: ARC1116

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 76 kDa

Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02%

Sodium Azide.


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




## Anti-PIAS1 Antibody [ARC1116] (A305639)

## Images:



Western blot analysis of extracts of various cell lines, using Anti-PIAS1 Antibody [ARC1116] (A305639) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 60s.



Western blot analysis of extracts of Mouse lung, using Anti-PIAS1 Antibody [ARC1116] (A305639) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Enhanced Kit (RM00021). Exposure time: 3min.