


Anti-IL-17A Antibody [ARC52762] (A306037)

Specifications:


| | |
|------------------------|--|
| Name: | Anti-IL-17A Antibody [ARC52762] |
| Description: | Rabbit monoclonal [ARC52762] antibody to IL-17A. |
| Applications: | IHC |
| Recommended Dilutions: | IHC: 1:50-1:200 |
| Reactivity: | Human, Mouse, Rat |
| Immunogen: | Recombinant fusion protein containing a sequence corresponding to amino acids 24-155 of human IL17A (NP_002181.1). |
| Sequence: | GITIPRNP GCPNSEDKNFPRTVMVNLNIHNRNTNTNPKRSSDY YNRSTSPWNLHRNED PERYPSVIWEAKCRHLGCINADGNVDYHMNSVPIQQEILVLRREPPHCPNSFRLEKIL VSVGCTCVTPIVHHVA |
| Host: | Rabbit |
| Clonality: | Monoclonal |
| Clone ID: | ARC52762 |
| Isotype: | IgG |
| Conjugate: | Unconjugated |
| Purification: | Affinity purification. |
| Product Form: | Liquid |
| Formulation: | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.05% Proclin 300. |
| Storage: | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |

Anti-IL-17A Antibody [ARC52762] (A306037)


Images:




Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue using Anti-IL-17A Antibody [ARC52762] (A306037) at a dilution of 1:100(40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human tonsil using Anti-IL-17A Antibody [ARC52762] (A306037) at a dilution of 1:100(40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse colon using Anti-IL-17A Antibody [ARC52762] (A306037) at a dilution of 1:100(40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat colon using Anti-IL-17A Antibody [ARC52762] (A306037) at a dilution of 1:100(40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.