


Anti-NF- κ B p65 Antibody [ARC51086] (A306573)

Specifications:


Name:	Anti-NF- κ B p65 Antibody [ARC51086]
Description:	Rabbit monoclonal [ARC51086] antibody to NF- κ B p65.
Applications:	WB, IHC, ICC/IF, IP, ChIP
Recommended Dilutions:	WB: 1:500-1:1,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200, IP: 1:50-1:200, ChIP: 1:50-1:200
Reactivity:	Human, Mouse, Rat, Monkey
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 450-551 of human NF- κ B p65 (Q04206).
Sequence:	LGALLGNSTDPVFTDLASVDNSEFQQLLNQGIPVAPHTTEPMLMEYPEAITRLVTGA QRPPDPAPAPLGAPGLPNGLLSGDEDFSSIADMDFSALLSQISS
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC51086
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	65 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-NF-kB p65 Antibody [ARC51086] (A306573)


Images:




Western blot analysis of extracts of various cell lines, using Anti-NF-kB p65 Antibody [ARC51086] (A306573) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 3min.



Western blot analysis of extracts of various cell lines, using Anti-NF-kB p65 Antibody [ARC51086] (A306573) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Enhanced Kit (RM00021). Exposure time: 3min.



Immunohistochemistry analysis of paraffin-embedded human colon tissue using Anti-NF-kB p65 Antibody [ARC51086] (A306573) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human tonsil using Anti-NF-kB p65 Antibody [ARC51086] (A306573) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.