


Anti-SQSTM1 / p62 Antibody [ARC0180] (A306736)

Specifications:


Name:	Anti-SQSTM1 / p62 Antibody [ARC0180]
Description:	Rabbit monoclonal [ARC0180] antibody to SQSTM1 / p62.
Applications:	WB, IHC, ICC/IF, IP
Recommended Dilutions:	WB: 1:500-1:1,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200, IP: 1:500-1:1,000
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 341-440 of human SQSTM1/p62 (Q13501).
Sequence:	LSSKEVDPSTGELQSLQMPESSEGPSSLDPSQEGPTGLKEAALYPHLPPEADPRLIESL SQMLSMGFSDDEGGWLRLLQTKNYDIGAALDTIQYSKHPPPL
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC0180
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	62 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-SQSTM1 / p62 Antibody [ARC0180] (A306736)


Images:




Western blot analysis of extracts of various cell lines, using Anti-SQSTM1 / p62 Antibody [ARC0180] (A306736) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.



Western blot analysis of extracts of PC-12 cells, using Anti-SQSTM1 / p62 Antibody [ARC0180] (A306736) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.



Western blot analysis of extracts of HeLa cells, using Anti-SQSTM1 / p62 Antibody [ARC0180] (A306736) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 180s.



Western blot analysis of extracts from wild type (WT) and SQSTM1/p62 knockout (KO) 293T cells, using Anti-SQSTM1 / p62 Antibody [ARC0180] (A306736) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.