


Anti-NRF1 Antibody [ARC0768] (A306763)

Specifications:


Name:	Anti-NRF1 Antibody [ARC0768]
Description:	Rabbit monoclonal [ARC0768] antibody to NRF1.
Applications:	WB, IHC, IP, ChIP
Recommended Dilutions:	WB: 1:500-1:2,000, IHC: 1:50-1:200, IP: 1:500-1:1,000, ChIP: 1:500-1:1,000
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 300-400 of human NRF1 (Q16656).
Sequence:	LVPSQTVVQTFSNPDGTVSLIQVGTGATVATLADASELPTTVTVAQVNYSAVADGEVE QNWATLQGGEMTIQTTQASEATQAVASLAEAAVAASQEMQQGA
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC0768
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	68 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-NRF1 Antibody [ARC0768] (A306763)


Images:




Western blot analysis of extracts of various cell lines, using Anti-NRF1 Antibody [ARC0768] (A306763) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.



Western blot analysis of extracts of various cell lines, using Anti-NRF1 Antibody [ARC0768] (A306763) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 5s.



Immunohistochemistry analysis of paraffin-embedded human colon tissue using Anti-NRF1 Antibody [ARC0768] (A306763) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunoprecipitation analysis of 300 μ g extracts of HeLa cells using 3 μ g of Anti-NRF1 Antibody [ARC0768] (A306763). This Western blot was performed on the immunoprecipitate using Anti-NRF1 Antibody [ARC0768] (A306763) at a dilution of 1:1000.