


## Anti-AHNAK Antibody (A307240)

### Specifications:

Name:	Anti-AHNAK Antibody
Description:	Rabbit polyclonal antibody to AHNAK.
Applications:	IHC, ICC/IF
Recommended Dilutions:	IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-149 of human AHNAK (NP_001611.1).
Sequence:	MEKEETTRELLLPNWQSGSHGLTIAQRDDGVFVQEVTQNSPAARTGVVKEGDQIVGA TIYFDNLQSGEVTQLLNTMGHHTVGLKLRKGD RSPEPGQWTREVFSSCSSEVVL SG DDEEYQRIYTTKIKPRLKSEDGVEGDLGETQSR
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


### Images:




Immunohistochemistry analysis of paraffin-embedded mouse liver using Anti-AHNAK Antibody (A307240) at a dilution of 1:50 (40x lens). 1. Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

## Anti-AHNAK Antibody (A307240)


Images continued:



Immunohistochemistry analysis of paraffin-embedded mouse brain using Anti-AHNAK Antibody (A307240) at a dilution of 1:50 (40x lens). 1. Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunofluorescence analysis of NIH/3T3 cells using Anti-AHNAK Antibody (A307240) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of A-549 cells using Anti-AHNAK Antibody (A307240) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).