


Anti-Cystatin-B Antibody [ARC2833] (A307531)

Specifications:


Name:	Anti-Cystatin-B Antibody [ARC2833]
Description:	Rabbit monoclonal [ARC2833] antibody to Cystatin-B.
Applications:	WB
Recommended Dilutions:	WB: 1:100-1:500
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 1-98 of human Cystatin B (P04080).
Sequence:	MMCGAPSATQPATAETQHIADQVRSQLEEKENKKFPVFKAVSFKSQVWAGTNYFIKVVH VGDEDFVHLRVFQSLPHENKPLTLSNYQTNKAKHDELTYF
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC2833
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	13 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-Cystatin-B Antibody [ARC2833] (A307531)

Images:



Western blot analysis of extracts of various cell lines, using Anti-Cystatin-B Antibody [ARC2833] (A307531) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.



Western blot analysis of extracts of various cell lines, using Anti-Cystatin-B Antibody [ARC2833] (A307531) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 180s.