


Anti-Ferritin Heavy Chain Antibody [ARC0023] (A307944)

Specifications:


Name:	Anti-Ferritin Heavy Chain Antibody [ARC0023]
Description:	Rabbit monoclonal [ARC0023] antibody to Ferritin Heavy Chain.
Applications:	WB
Recommended Dilutions:	WB: 1:500-1:2,000
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Ferritin Heavy Chain (P02794).
Sequence:	MTTASTSQVRQNYHQDSEAAINRQINLELYASYVYLSMSYYFDRDDVALKNFAKYFLH QSHEEREHAEKLMKLNQRGGRIFLQDIKKPDCDDWESGLNA
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC0023
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	21 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-Ferritin Heavy Chain Antibody [ARC0023] (A307944)

Images:



Western blot analysis of extracts of U-251MG cells, using Anti-Ferritin Heavy Chain Antibody [ARC0023] (A307944) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 3min.



Western blot analysis of extracts of various cell lines, using Anti-Ferritin Heavy Chain Antibody [ARC0023] (A307944) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.