


Anti-USP22 Antibody [ARC1498] (A307989)

Specifications:


Name:	Anti-USP22 Antibody [ARC1498]
Description:	Rabbit monoclonal [ARC1498] antibody to USP22.
Applications:	WB
Recommended Dilutions:	WB: 1:500-1:2,000
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-175 of human USP22 (Q9UPT9).
Sequence:	MVSRPEPEGEAMDAELAVAPPGCSHLGSFKVDNWKQNLRAIYQCFVWSGTAEARKRKA KSCICHVCGVHNLRLHSLCYVFFGCFSTKKHIHEHAKAKRHNLAIDLMYGGIYCFLCQ DYIYDKDMEIIAKEEQRKAWKMQGVGEEKFSTWEPTKRELELLKHNPKRRKITSNCTIG L
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC1498
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	60 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-USP22 Antibody [ARC1498] (A307989)


Images:



Western blot analysis of extracts of various cell lines, using Anti-USP22 Antibody [ARC1498] (A307989) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.



Western blot analysis of extracts of Mouse thymus, using Anti-USP22 Antibody [ARC1498] (A307989) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.



Western blot analysis of extracts of Rat lung, using Anti-USP22 Antibody [ARC1498] (A307989) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 3min.