

## Anti-KAP1 Antibody [ARC0047] (A308504)

## Specifications:

Name: Anti-KAP1 Antibody [ARC0047]

Description: Rabbit monoclonal [ARC0047] antibody to KAP1.

Applications: WB, IHC, IP

Recommended Dilutions: WB: 1:500-1:2,000, IHC: 1:50-1:200, IP: 1:500-1:1,000

Reactivity: Human, Mouse, Rat

Immunogen: A synthetic peptide corresponding to a sequence within amino acids 250-350 of human

KAP1/TRIM28 (Q13263).

Sequence: RNQRKLLASLVKRLGDKHATLQKSTKEVRSSIRQVSDVQKRVQVDVKMAILQIMKELN

KRGRVLVNDAQKVTEGQQERLERQHWTMTKIQKHQEHILRFAS

Host: Rabbit

Clonality: Monoclonal

Clone ID: ARC0047

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 110 kDa

Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02%

Sodium Azide.


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




## Anti-KAP1 Antibody [ARC0047] (A308504)


## Images:




Western blot analysis of extracts of various cell lines, using Anti-KAP1 Antibody [ARC0047] (A308504) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit lgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.



Immunohistochemistry analysis of paraffin-embedded rat spleen using Anti-KAP1 Antibody [ARC0047] (A308504) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human appendix tissue using Anti-KAP1 Antibody [ARC0047] (A308504) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse brain using Anti-KAP1 Antibody [ARC0047] (A308504) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.