


## Anti-RBMY1A1 Antibody (A308659)

### Specifications:

Name:	Anti-RBMY1A1 Antibody
Description:	Rabbit polyclonal antibody to RBMY1A1.
Applications:	IHC, ICC/IF
Recommended Dilutions:	IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 90-300 of human RBMY1A1 (NP_005049.1).
Sequence:	SGGRRRPPASSRNRSPSGSLRSARGSRGGTRGWLPSEHGHLDDGGYTPDLKMSYSRGL IPVKRGPSSRSGGPPPKKSAPSAVARNSWMGSQGPMSQRRENYGVPPRRATISSWRN DRMSTRHDGYATNDGNHPSCQETRDIYAPPSRGYAYRDNGHSNRDEHSSRGYRNHRSSR ETRDYAPPSRGHAYRDYGHSSRRDESYSRGYRNRRSSR
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


### Images:




Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue using Anti-RBMY1A1 Antibody (A308659) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

## Anti-RBMY1A1 Antibody (A308659)


Images continued:



Immunohistochemistry analysis of paraffin-embedded mouse lung using Anti-RBMY1A1 Antibody (A308659) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat kidney using Anti-RBMY1A1 Antibody (A308659) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunofluorescence analysis of mouse testis cells using Anti-RBMY1A1 Antibody (A308659) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).