


Anti-Doublecortin Antibody (A308716)

Specifications:

Name:	Anti-Doublecortin Antibody
Description:	Rabbit polyclonal antibody to Doublecortin.
Applications:	WB, ICC/IF
Recommended Dilutions:	WB: 1:100-1:500, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 265-365 of human Doublecortin (NP_835365.1).
Sequence:	SLDENECRVMKGNPSATAGPKASPTPQKTSKSPGPMRRSKSPADSGNDQDANGTSSS QLSTPKSKQSPISTPTSPGSLRKHKDLYLPLSLDDSDSLGDSM
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	41 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:




Western blot analysis of extracts of various cell lines, using Anti-Doublecortin Antibody (A308716) at 1:400 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.

Anti-Doublecortin Antibody (A308716)


Images continued:



Immunofluorescence analysis of SH-SY5Y cells using Anti-Doublecortin Antibody (A308716) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of rat brain using Anti-Doublecortin Antibody (A308716) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of mouse brain using Anti-Doublecortin Antibody (A308716) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).