
antibodies

Anti-CTLA4 Antibody [ARC57390] (A309628)

Specifications:

| Name: | Anti-CTLA4 Antibody [ARC57390] |
|------------------------|---|
| Description: | Rabbit monoclonal [ARC57390] antibody to CTLA4. |
| Applications: | WB |
| Recommended Dilutions: | WB: 1:2,000-1:10,000 |
| Reactivity: | Human |
| Immunogen: | Recombinant protein of human CD152/CTLA-4. |
| Host: | Rabbit |
| Clonality: | Monoclonal |
| Clone ID: | ARC57390 |
| Isotype: | lgG |
| Conjugate: | Unconjugated |
| Purification: | Affinity purification. |
| Molecular Weight: | 25 kDa |
| Product Form: | Liquid |
| Formulation: | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.05% Proclin 300. |
| Storage: | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |

Images:




Western blot analysis of 293F, using Anti-CTLA4 Antibody [ARC57390] (A309628) at 1:10,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 60s.

antibodies

Anti-CTLA4 Antibody [ARC57390] (A309628)

Images continued:



Western blot analysis of Recombinant Human CTLA-4 Protein, using Anti-CTLA4 Antibody [ARC57390] (A309628) at 1:10,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 60s.