

Anti-CD40 Antibody [DM101] - BSA and Azide free (A318604)

Specifications:

Name: Anti-CD40 Antibody [DM101] - BSA and Azide free

Description: Recombinant rabbit monoclonal [DM101] antibody to CD40.

Applications: ELISA, Flow Cytometry

Recommended Dilutions: ELISA: 1:5,000-10,000, Flow Cytometry: 1:100

Reactivity: Human

Host: Rabbit

Clonality: Monoclonal

Clone ID: DM101

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity chromatography.

Concentration: Reconstitution dependent.

Product Form: Lyophilized

Reconstitution: Reconstitute with distilled sterile water.

Formulation: Lyophilized from sterile Phosphate Buffered Saline, pH 7.4. Normally 5%-8% Trehalose is

added as a protectant before lyophilization.

Storage: Shipped at 4°C. Lyophilized: Store at -20°C to -80°C. Reconstituted: Aliquot and store at

-80°C. Product is stable for one year. Avoid freeze/thaw cycles.

General Notes: Prior to reconstitution, centrifuge the vial at 5,000g for 3-5 minutes at room temperature.

Reconstitute with appropriate volume of distilled sterile water to bring product to 1mg/ml concentration. After addition of distilled sterile water, mix by gentle tapping. Note: It is not


recommended to vortex or vigorously pipette the sample.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




Anti-CD40 Antibody [DM101] - BSA and Azide free (A318604)


Images:



ELISA plates were pre-coated with Recombinant Human CD40 Protein (Fc Chimera 6xHis Tag) (A318393) at 2 μ g/ml (100 μ l/well) which can bind Anti-CD40 Antibody [DM101] - BSA and Azide free (A318604) in a linear range of 0.64-80 ng/ml.



A. Flow cytometry analysis with Anti-CD40 Antibody [DM101] - BSA and Azide free (A318604) on Expi293 cells transfected with human CD40 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).



Flow cytometry analysis of antigen binding of Anti-CD40 Antibody [DM101] - BSA and Azide free (A318604). (A) Anti-CD40 Antibody [DM101] - BSA and Azide free (A318604) does not bind to Jurkat cells that do not express CD40. (B) A clear peak shift of Anti-CD40 Antibody [DM101] - BSA and Azide free (A318604) was seen compared to the control when incubated with CD40-expressing Raji cells, indicating strong binding of Anti-CD40 Antibody [DM101] - BSA and Azide free (A318604) to CD40. Antibodies were incubated at 2 μ g/ml.