

Etigilimab Biosimilar - Anti-TIGIT Antibody - BSA and Azide free (A318932)


Specifications:

Name:	Etigilimab Biosimilar - Anti-TIGIT Antibody - BSA and Azide free
Description:	Recombinant humanized monoclonal antibody to TIGIT.
Applications:	ELISA, Flow Cytometry
Recommended Dilutions:	ELISA: 1:5,000-10,000, Flow Cytometry: 1:100
Reactivity:	Human
Host:	Humanized
Clonality:	Monoclonal
lsotype:	lgG1
Conjugate:	Unconjugated
Purification:	Affinity chromatography.
Concentration:	Reconstitution dependent.
Product Form:	Lyophilized
Reconstitution:	Reconstitute with distilled sterile water.
Formulation:	Lyophilized from sterile Phosphate Buffered Saline, pH 7.4. Normally 5%-8% Trehalose is added as a protectant before lyophilization.
Storage:	Shipped at 4°C. Lyophilized: Store at -20°C to -80°C. Reconstituted: Aliquot and store at -80°C. Product is stable for one year. Avoid freeze/thaw cycles.
General Notes:	Prior to reconstitution, centrifuge the vial at 5,000g for 3-5 minutes at room temperature. Reconstitute with appropriate volume of distilled sterile water to bring product to 1mg/ml concentration. After addition of distilled sterile water, mix by gentle tapping. Note: It is not recommended to vortex or vigorously pipette the sample.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.



Etigilimab Biosimilar - Anti-TIGIT Antibody - BSA and Azide free (A318932)

Images:




ELISA plates were pre-coated with Recombinant Human TIGIT Protein (6×His Tag) (A318229) at 2 µg/ml (100 µl/well) which can bind Etigilimab Biosimilar - Anti-TIGIT Antibody - BSA and Azide free (A318932) in a linear range of 0.13-16.0 ng/ml.

Anti-TIGIT (etigilimab biosimilar) mAb ELISA fean Abs.(OD450) 100 Anti-TIGIT Neutralizing a

body(ng/ml)

ELISA plates were pre-coated with Recombinant Human TIGIT Protein (Fc Tag) (A318230) at 2 µg/ml (100 µl/well) which can bind Etigilimab Biosimilar - Anti-TIGIT Antibody - BSA and Azide free (A318932) in a linear range of 0.13-16.0 ng/ml.



HEK293 cell line was transfected with: Irrelevant protein (B) and human TIGIT protein (A). Cells were surface stained with 1µg/ml of Etigilimab Biosimilar - Anti-TIGIT Antibody - BSA and Azide free (A318932) followed by Anti-Human IgG Antibody (Alexa 488).