


Anti-IgG Antibody [I17-T] (A8243)

Specifications:

Description:Rabbit monoclonal (I17-T) antibody to IgG.Specificity:This antibody recognizes the epitope between Ala261 - Lys275.Applications:IHC-P, IHC-FrRecommended Dilutions:IHC: 1:100-1:300.Reactivity:HumanImmunogen:Peptide derived from the domain close to the C-terminus of human IgG-1 chain C region.	Name:	Anti-IgG Antibody [I17-T]
Applications:IHC-P, IHC-FrRecommended Dilutions:IHC: 1:100-1:300.Reactivity:HumanImmunogen:Peptide derived from the domain close to the C-terminus of human IgG-1 chain C region.	Description:	Rabbit monoclonal (I17-T) antibody to IgG.
Recommended Dilutions:IHC: 1:100-1:300.Reactivity:HumanImmunogen:Peptide derived from the domain close to the C-terminus of human IgG-1 chain C region.	Specificity:	This antibody recognizes the epitope between Ala261 - Lys275.
Reactivity:HumanImmunogen:Peptide derived from the domain close to the C-terminus of human IgG-1 chain C region.	Applications:	IHC-P, IHC-Fr
Immunogen: Peptide derived from the domain close to the C-terminus of human IgG-1 chain C region.	Recommended Dilutions:	IHC: 1:100-1:300.
	Reactivity:	Human
Debb'	Immunogen:	Peptide derived from the domain close to the C-terminus of human IgG-1 chain C region.
HOST: Kaddit	Host:	Rabbit
Clonality: Monoclonal	Clonality:	Monoclonal
Clone ID: I17-T	Clone ID:	I17-T
Isotype: IgG	lsotype:	IgG
Conjugate: Unconjugated	Conjugate:	Unconjugated
Product Form: Liquid	Product Form:	Liquid
Formulation: Supplied in Tris Buffered Saline, pH 8.0, with 20 mg/ml BSA and 0.05% Sodium Azide.	Formulation:	Supplied in Tris Buffered Saline, pH 8.0, with 20 mg/ml BSA and 0.05% Sodium Azide.
Storage: Store at 4°C.	Storage:	Store at 4°C.
Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.	Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Images:



lgG positive plasma cells in the case of lgG4-related chronic sclerosing sialoadenitis, detected with Anti-IgG Antibody (A8243). Formalin fixed, paraffin embedded human tissue (4 μ m section) stained.