


## Anti-p53 Antibody (A80444)

### Specifications:


Name:	Anti-p53 Antibody
Description:	Rabbit polyclonal antibody to p53.
Applications:	IHC
Recommended Dilutions:	IHC: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 95-390 of mouse p53 (NP_035770.2).
Sequence:	PSQKTYQGNYGFHLGFLQSGTAKSVMCTYSPPLNKLFCQLAKTCPVQLWVSATPPAGS RVRAMAIYKKSQHMTEVRRRCPPHHERCSDGDGLAPPQHLIRVEGNLYPEYLEDQRQTFR HSVVVPYEPPEAGSEYTTIHYKYMCSMGGMNRRPILTIITLEDSSGNLLGRDSFE VRVCACPGRRRTEENFRKKEVLCPELPPGSAKRALPTCTSASPPQKKKPLDGEYFT LKIRGRKRFEMFRELNEALELKDAHATEESGDSRAHSSYLKTKKGQSTSRHKKTMVKK VGPDS
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-p53 Antibody (A80444)


### Images:




Immunohistochemistry analysis of paraffin-embedded rat brain using Anti-p53 Antibody (A80444) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human lung cancer using Anti-p53 Antibody (A80444) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue using Anti-p53 Antibody (A80444) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse lung using Anti-p53 Antibody (A80444) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.