


## Anti-Stromal interaction molecule 1 Antibody (A87712)

### Specifications:


Name:	Anti-Stromal interaction molecule 1 Antibody
Description:	Rabbit polyclonal antibody to Stromal interaction molecule 1.
Applications:	WB, IHC, ICC/IF, IP
Recommended Dilutions:	WB: 1:500-1:1,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200, IP: 1:500-1:1,000
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 530-685 of human STIM1 (NP_003147.2).
Sequence:	RQRVAPKPPQMSRAADEALNAMTSNGSHRLIEGVHPGSLVEKLPDSPALAKKALLALN HGLDKAHSMLSPSAPPGGSPHLDSSRSHPSSPDPDTPSPVGDSTRALQASRNTRIP HLAGKKAVAEEDNGSIGEETDSSPGRKKFPLKIFKKPLKK
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	85 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-Stromal interaction molecule 1 Antibody (A87712)


### Images:




Western blot analysis of extracts of various cell lines, using Anti-Stromal interaction molecule 1 Antibody (A87712) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.



Immunofluorescence analysis of HeLa cells using Anti-Stromal interaction molecule 1 Antibody (A87712) at a dilution of 1:50 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of NIH/3T3 cells using Anti-Stromal interaction molecule 1 Antibody (A87712) at a dilution of 1:50 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunoprecipitation analysis of 300µg extracts of HeLa cells using 3µg of Anti-Stromal interaction molecule 1 Antibody (A87712). This Western blot was performed on the immunoprecipitate using Anti-Stromal interaction molecule 1 Antibody (A87712) at a dilution of 1:1000.