


## Anti-SCD1 Antibody (A89598)

### Specifications:


Name:	Anti-SCD1 Antibody
Description:	Rabbit polyclonal antibody to SCD1.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:500-1:1,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 200 to the C-terminus of human SCD (NP_005054.3).
Sequence:	LDLSDLEAEKLVMFQRRYYKPGLLMMCFILPTLVPWYFWGETFQNSVVFVATFLRYAVV LNATWLVNSAAHLFGYRYPYDKNISPRENILVSLGAVGEGFHNYHHSFPYDYSASEYRW HINFITTFIDCMAALGLAYDRKKVSKAAILARIKRTGDGNYKSG
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	37 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-SCD1 Antibody (A89598)


### Images:




Western blot analysis of extracts of various cell lines, using Anti-SCD1 Antibody (A89598) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.



Western blot analysis of extracts of U-251MG cells, using Anti-SCD1 Antibody (A89598) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 3s.



Immunohistochemistry analysis of paraffin-embedded rat brain using Anti-SCD1 Antibody (A89598) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse brain using Anti-SCD1 Antibody (A89598) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.